

## පැරණි නිර්දේශය/பழைய பாடத்திட்டம்/Old Syllabus

ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව  
 திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம்  
 Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka  
 இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம்  
 Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka

OLD

රසායන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 2020  
 கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2020  
 General Certificate of Education (Adv. Level) Examination, 2020

රසායන විද්‍යාව I  
 இரசாயனவியல் I  
 Chemistry I

02 E I

පැය දෙකයි  
 இரண்டு மணித்தியாலம்  
 Two hours

## Instructions:

- \* Periodic Table is provided.
- \* This paper consists of 09 pages.
- \* Answer all the questions.
- \* Use of calculators is not allowed.
- \* Write your Index Number in the space provided in the answer sheet.
- \* Follow the instructions given on the back of the answer sheet carefully.
- \* In each of the questions 1 to 50, pick one of the alternatives from (1), (2), (3), (4), (5) which is correct or most appropriate and mark your response on the answer sheet with a cross (x) in accordance with the instructions given on the back of the answer sheet.

Universal gas constant  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$   
 Avogadro constant  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Planck's constant  $h = 6.626 \times 10^{-34} \text{ Js}$   
 Velocity of light  $c = 3 \times 10^8 \text{ ms}^{-1}$

1. Identify the incorrect statement from the following.

- (1) The filling of electrons into orbitals of equal energy is governed by Hund's rule.
- (2) Wave nature of electrons has been shown by diffraction experiments.
- (3) In hydrogen, when electrons fall from high energy levels to the level with principal quantum number,  $n = 1$ , the line spectrum observed is called Lyman series.
- (4) Atoms absorb or emit radiation in the form of definite small quantities and the smallest quantity is referred to as a photon.
- (5) Two electrons in an orbital must have opposite spins as deduced from Aufbau principle.

2. The number of electrons in the manganese atom (Mn,  $Z = 25$ ) that have quantum numbers  $l = 0$  and  $m_l = -1$  respectively are,

- (1) 6 and 4      (2) 8 and 12      (3) 8 and 5      (4) 8 and 6      (5) 10 and 5

3. M is an element that belongs to the second period in the Periodic Table. It forms a covalent molecule  $MCl_3$  which has a dipole moment. The group of the Periodic Table to which M belongs is,

- (1) 2      (2) 13      (3) 14      (4) 15      (5) 16


4. The number of unstable Lewis structures that can be drawn for the peroxyntic acid molecule

(formula  $\text{HNO}_4$ ,  $\text{H}-\ddot{\text{O}}-\ddot{\text{O}}-\overset{\text{:O:}}{\underset{\oplus}{\text{N}}}-\ddot{\text{O}}^-$ ) is,

- (1) 1      (2) 2      (3) 3      (4) 4      (5) 5

5. The IUPAC name of the given compound is,

- (1) 1-bromo-4-methyl-5-hydroxypent-1-en-3-one
- (2) 5-bromo-1-hydroxy-2-methylpent-4-en-3-one
- (3) 1-bromo-5-hydroxy-4-methylpent-1-en-3-one
- (4) 5-bromo-2-methyl-3-oxopent-4-en-1-ol
- (5) 1-bromo-4-methyl-3-oxopent-1-enol



6. The decreasing order of radii of the species O, O<sup>2-</sup>, F, F<sup>-</sup>, S<sup>2-</sup>, Cl<sup>-</sup> is,

- (1) S<sup>2-</sup> > Cl<sup>-</sup> > O<sup>2-</sup> > F<sup>-</sup> > O > F  
 (2) S<sup>2-</sup> > Cl<sup>-</sup> > O<sup>2-</sup> > F<sup>-</sup> > F > O  
 (3) Cl<sup>-</sup> > S<sup>2-</sup> > O<sup>2-</sup> > F<sup>-</sup> > O > F  
 (4) Cl<sup>-</sup> > S<sup>2-</sup> > F<sup>-</sup> > O<sup>2-</sup> > O > F  
 (5) S<sup>2-</sup> > Cl<sup>-</sup> > O<sup>2-</sup> > O > F > F

7. A rigid-closed container contains  $n_1$  moles of an ideal gas at temperature  $T_1$ (K) and pressure  $P_1$ (Pa). When an additional amount of the gas was inserted into the container, the new temperature and pressure were  $T_2$  and  $P_2$ , respectively. The total number of moles of the gas now in the container is,

- (1)  $\frac{n_1 T_1 P_1}{T_2 P_2}$  (2)  $\frac{n_1 T_1 P_1}{T_2 P_1}$  (3)  $\frac{T_2 P_2}{n_1 T_1 P_1}$  (4)  $\frac{n_1 T_1 P_2}{T_1 P_1}$  (5)  $\frac{n_1 T_1 P_1}{T_1 P_2}$

8. The total number of electrons exchanged in the reaction of the oxidation of ethanol (C<sub>2</sub>H<sub>5</sub>OH) to acetic acid (CH<sub>3</sub>COOH) using acidic K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> solution is,

- (1) 6 (2) 8 (3) 10 (4) 12 (5) 14

9. Which compound of the following, can undergo aldol condensation, when reacted with aqueous NaOH?

- (1) CH<sub>3</sub>C(=O)OH (2) CH<sub>3</sub>C(=O)OCH<sub>3</sub> (3) H-C(=O)OCH<sub>3</sub> (4) CH<sub>3</sub>CH<sub>2</sub>C(=O)H (5) (CH<sub>3</sub>)<sub>3</sub>CC(=O)H


10. AX(s), A<sub>2</sub>Y(s) and AZ(s) are sparingly soluble salts in water having  $K_{sp}$  values of  $1.6 \times 10^{-9}$ ,  $3.2 \times 10^{-11}$  and  $9.0 \times 10^{-12}$ , respectively at 25 °C. Which of the following shows the order of the three saturated solutions of these salts in decreasing concentration of cation A<sup>+</sup>(aq), at 25 °C ?

- (1) AX(s) > A<sub>2</sub>Y(s) > AZ(s)  
 (2) A<sub>2</sub>Y(s) > AX(s) > AZ(s)  
 (3) AX(s) > AZ(s) > A<sub>2</sub>Y(s)  
 (4) A<sub>2</sub>Y(s) > AZ(s) > AX(s)  
 (5) AZ(s) > A<sub>2</sub>Y(s) > AX(s)

11. Consider the following compounds.

	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	$\begin{array}{c} \text{CH}_3 \\   \\ \text{CH}_3\text{CCH}_2\text{CH}_3 \\   \\ \text{CH}_3 \end{array}$	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CHO	$\begin{array}{c} \text{CH}_3 \\   \\ \text{CH}_3\text{CCHO} \\   \\ \text{CH}_3 \end{array}$	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH
	A	B	C	D	E
Relative molecular mass	86	86	86	86	88

Variation of boiling points of these compounds is best shown by,




12. The increasing order of covalent character of the chemical species NaCl, Na<sub>2</sub>S, KF and KCl is,

- (1) KF < NaCl < KCl < Na<sub>2</sub>S  
 (2) KCl < NaCl < KF < Na<sub>2</sub>S  
 (3) KF < KCl < NaCl < Na<sub>2</sub>S  
 (4) Na<sub>2</sub>S < NaCl < KCl < KF  
 (5) KF < Na<sub>2</sub>S < NaCl < KCl

13. Standard combustion enthalpies of H<sub>2</sub>(g), C(s) and CH<sub>3</sub>OH(l) at 298 K are -286 kJ mol<sup>-1</sup>, -393 kJ mol<sup>-1</sup> and -726 kJ mol<sup>-1</sup>, respectively. Enthalpy of vaporization of CH<sub>3</sub>OH(l) is +37 kJ mol<sup>-1</sup>. Enthalpy of formation (kJ mol<sup>-1</sup>) of one mole of gaseous CH<sub>3</sub>OH at 298 K is,

- (1) -276                      (2) -239                      (3) -202                      (4) +84                      (5) +202


14. Phosphorous can be prepared in an electric furnace as given by the following balanced chemical equation.



When 620 g of Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>, 180 g of SiO<sub>2</sub> and 96 g of C were reacted, 50 g of P<sub>4</sub> were obtained. Under these conditions, the limiting reagent (reagent that is completely consumed) and percentage yield of P<sub>4</sub> respectively are, (C = 12, O = 16, Si = 28, P = 31, Ca = 40)

- (1) Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> and 80.7%                      (2) SiO<sub>2</sub> and 80.7%                      (3) C and 50.4%  
 (4) SiO<sub>2</sub> and 40.3%                      (5) C and 25.2%






15. Consider the following two equilibria occurring in two separate rigid-closed containers under the same conditions.



Under these conditions K<sub>p</sub> for the equilibrium 2H<sub>2</sub>S(g) + N<sub>2</sub>(g) + 3H<sub>2</sub>(g) ⇌ 2NH<sub>4</sub>HS(g) is,


- (1) 5.76 × 10<sup>-12</sup>                      (2) 7.2 × 10<sup>-10</sup>                      (3) 1.92 × 10<sup>-8</sup>                      (4) 3.40 × 10<sup>-6</sup>                      (5) 3.75 × 10<sup>-2</sup>

16. Consider the nitration reaction of bromobenzene. Resonance stabilized carbocation intermediates are formed during this reaction. Which of the following is **not** a resonance structure of these intermediates?




17. A reaction which is non-spontaneous at room temperature and 1 atm pressure becomes spontaneous at high temperature at the same pressure. Which of the following is correct for this reaction at room temperature? (Assume that ΔH and ΔS do not change with temperature and pressure.)

- |     | ΔG       | ΔH       | ΔS       |
|-----|----------|----------|----------|
| (1) | Positive | Positive | Positive |
| (2) | Positive | Negative | Negative |
| (3) | Positive | Negative | Positive |
| (4) | Negative | Positive | Negative |
| (5) | Negative | Negative | Negative |

18. The charges on the central sulphur and oxygen atoms in the most acceptable Lewis structure for the SO<sub>4</sub> molecule (skeleton: ) respectively are,

- (1) +1 and zero                      (2) zero and -1                      (3) zero and zero  
 (4) +2 and zero                      (5) +2 and -1


19. Which of the following correctly shows the electrolytic cell constructed for the electrolysis of an aqueous solution of the salt MX?



20. Which of the following statements is correct regarding the reaction between a carboxylic acid and an alcohol to give an ester?

- (1) The overall reaction is a nucleophilic addition reaction of a carbonyl compound.
- (2) It is a reaction in which the alcohol acts as a nucleophile.
- (3) It is a reaction which occurs with the cleavage of the O–H bond of the carboxylic acid.
- (4) It is a reaction which occurs with the cleavage of the C–O bond of the alcohol.
- (5) It is an acid-base reaction.

21. Decomposition of 1 mol of  $\text{CH}_3\text{OH}(l)$  occurs at high temperatures as follows.



Which of the following is **incorrect** for the above reaction? (H = 1, C = 12, O = 16)

- (1) The heat absorbed when 1 mol of  $\text{CH}_3\text{OH}(g)$  is decomposed is less than 128 kJ.
- (2) Enthalpy of  $\text{CO}(g) + 2\text{H}_2(g)$  is higher than the enthalpy of  $\text{CH}_3\text{OH}(l)$ .
- (3) 128 kJ of heat is released when 1 mol of  $\text{CO}(g)$  is formed.
- (4) 128 kJ of heat is absorbed during the decomposition of a mole of reactant.
- (5) 128 kJ of heat is absorbed when 32 g of products are formed.

22. Identify the **incorrect** statement from the following.

- (1) Electron affinity of nitrogen  $[\text{N}(g)]$  is positive.
- (2) Dilution of  $\text{BiCl}_3(\text{aq})$  solution with water gives a white precipitate.
- (3)  $\text{H}_2\text{S}$  gas can act both as an oxidizing agent and a reducing agent.
- (4) The effective nuclear charge ( $Z^*$ ) felt by a valence electron in He is less than 2.
- (5) When sulphur reacts with conc.  $\text{H}_2\text{SO}_4$ ,  $\text{SO}_3$  gas is produced

23. The concentration of a dilute aqueous solution of a weak acid HA is  $C \text{ mol dm}^{-3}$  and its acid dissociation constant is  $K_a$  at 298 K. Which of the following expressions gives the pH of the solution at 298 K?

- (1)  $\text{pH} = \frac{1}{2} \text{p}K_a - \frac{1}{2} \log C$
- (2)  $\text{pH} = -\frac{1}{2} \text{p}K_a - \frac{1}{2} \log C$
- (3)  $\text{pH} = -\frac{1}{2} \text{p}K_a + \frac{1}{2} \log C$
- (4)  $\text{pH} = -\frac{1}{2} \text{p}K_a - \frac{1}{2} \log (1/C)$
- (5)  $\text{pH} = \frac{1}{2} \text{p}K_a - \frac{1}{2} \log (1/C)$

24. The strength of a  $\text{H}_2\text{O}_2$  solution can be expressed as the volume of  $\text{O}_2$  produced at standard temperature and pressure (STP). For example, a litre of 20 **volume strength**  $\text{H}_2\text{O}_2$  solution will produce 20 litres of  $\text{O}_2$  gas at STP ( $2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$ ). (Assume that 1 mole of gas has 22.4 litres volume at STP.)

A bottle labelled X contains  $\text{H}_2\text{O}_2$  solution. When  $25.0 \text{ cm}^3$  of solution X was titrated with  $1.0 \text{ mol dm}^{-3}$   $\text{KMnO}_4$  in the presence of dilute  $\text{H}_2\text{SO}_4$  the volume required to reach the end point was  $25.0 \text{ cm}^3$ . The volume strength of solution X is,

- (1) 15                      (2) 20                      (3) 25                      (4) 28                      (5) 30

25.  $\text{M}(\text{OH})_2(\text{s})$  is a sparingly water soluble salt formed by the reaction between  $\text{M}^{2+}(\text{aq})$  and  $\text{OH}^-(\text{aq})$  ions at 298 K. The solubility ( $\text{mol dm}^{-3}$ ) of  $\text{M}(\text{OH})_2(\text{s})$  in water at  $\text{pH} = 5$  is, ( $K_{sp}\text{M}(\text{OH})_2 = 4.0 \times 10^{-36}$  at 298 K).

- (1)  $\sqrt{2} \times 10^{-18}$       (2)  $2 \times 10^{-18}$       (3)  $1 \times 10^{-18}$       (4)  $\sqrt[3]{2} \times 10^{-12}$       (5)  $1 \times 10^{-12}$

26. Which of the following correctly denotes the standard galvanic cell constructed by using a standard hydrogen electrode, a standard Mg-electrode and a salt-bridge at 298 K?

- (1)  $\text{Mg}(\text{s}) \mid \text{Mg}^{2+}(\text{aq}, 1.00 \text{ mol dm}^{-3}) \parallel \text{H}^+(\text{aq}, 1.00 \text{ mol dm}^{-3}) \mid \text{H}_2(\text{g}) \mid \text{Pt}(\text{s})$   
 (2)  $\text{Pt}(\text{s}) \mid \text{H}_2(\text{g}) \mid \text{H}^+(\text{aq}, 1.00 \text{ mol dm}^{-3}) \parallel \text{Mg}^{2+}(\text{aq}, 1.00 \text{ mol dm}^{-3}) \mid \text{Mg}(\text{s})$   
 (3)  $\text{Mg}(\text{s}), \text{Mg}^{2+}(\text{aq}, 1.00 \text{ mol dm}^{-3}) \parallel \text{H}^+(\text{aq}, 1.00 \text{ mol dm}^{-3}) \mid \text{H}_2(\text{g}) \mid \text{Pt}(\text{s})$   
 (4)  $\text{Mg}(\text{s}) \mid \text{Mg}^{2+}(\text{aq}, 1.00 \text{ mol dm}^{-3}), \text{H}^+(\text{aq}, 1.00 \text{ mol dm}^{-3}), \text{H}_2(\text{g}) \mid \text{Pt}(\text{s})$   
 (5)  $\text{Pt}(\text{s}), \text{H}_2(\text{g}) \mid \text{H}^+(\text{aq}, 1.00 \text{ mol dm}^{-3}) \parallel \text{Mg}^{2+}(\text{aq}, 1.00 \text{ mol dm}^{-3}), \text{Mg}(\text{s})$


27. The following procedure was carried out at 298 K to determine the distribution coefficient  $K_D$  of a monobasic organic acid between dichloromethane and water.  $50.00 \text{ cm}^3$  of a  $0.20 \text{ mol dm}^{-3}$  aqueous solution of acid were mixed vigorously with  $10.00 \text{ cm}^3$  of dichloromethane and the two layers were allowed to separate. Thereafter, the dichloromethane layer in the bottom of the flask was drained out.  $10.00 \text{ cm}^3$  of  $0.02 \text{ mol dm}^{-3}$   $\text{NaOH}(\text{aq})$  solution were required to neutralize the acid remaining in the aqueous layer. (Assume that the acid does not dimerize in the organic phase.)  $K_D$  of the acid between **dichloromethane and water** at 298 K is,

- (1) 0.05                      (2) 0.25                      (3) 4.00                      (4) 20.00                      (5) 245.00

28. A reaction  $\text{C}_2\text{H}_4(\text{g}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g})$  occurs in a rigid-closed container at a given temperature. After a certain time, it was found that the rate of the reaction with respect to consumption of  $\text{C}_2\text{H}_4(\text{g})$  was  $x \text{ mol dm}^{-3} \text{ s}^{-1}$ . Which of the following shows the rates of consumption of  $\text{O}_2(\text{g})$ , formation of  $\text{CO}_2(\text{g})$  and formation of  $\text{H}_2\text{O}(\text{g})$  respectively, during that time?

	rate / $\text{mol dm}^{-3} \text{ s}^{-1}$		
	$\text{O}_2(\text{g})$	$\text{CO}_2(\text{g})$	$\text{H}_2\text{O}(\text{g})$
(1)	$\frac{3}{x}$	$\frac{2}{x}$	$\frac{2}{x}$
(2)	$x$	$x$	$x$
(3)	$\frac{x}{3}$	$\frac{x}{2}$	$\frac{x}{2}$
(4)	$\frac{1}{x}$	$\frac{1}{x}$	$\frac{1}{x}$
(5)	$3x$	$2x$	$2x$


29. Consider the following reaction occurring in a rigid-closed container at temperature  $T$ .



The rate of reaction doubled when the concentration of M was doubled. The rate of reaction is  $5.00 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$  when the concentrations of M and Q are  $1.0 \times 10^{-5} \text{ mol dm}^{-3}$  and  $2.0 \text{ mol dm}^{-3}$  respectively. The rate constant of the reaction under these conditions is,

- (1)  $2.5 \times 10^{-4} \text{ s}^{-1}$       (2)  $12.5 \text{ s}^{-1}$       (3)  $25 \text{ s}^{-1}$       (4)  $50 \text{ s}^{-1}$       (5)  $500 \text{ s}^{-1}$

30. Consider the following reaction scheme.



● For each of the questions 31 to 40, one or more responses out of the four responses (a), (b), (c) and (d) given is/are correct. Select the correct response/responses. In accordance with the instructions given on your answer sheet, mark

- (1) if only (a) and (b) are correct.  
 (2) if only (b) and (c) are correct.  
 (3) if only (c) and (d) are correct.  
 (4) if only (d) and (a) are correct.  
 (5) if any other number or combination of responses is correct.


**Summary of above Instructions**

(1)	(2)	(3)	(4)	(5)
Only (a) and (b) are correct	Only (b) and (c) are correct	Only (c) and (d) are correct	Only (d) and (a) are correct	Any other number or combination of responses is correct

31. Which of the following statement/s is/are correct with regard to 3d-block elements and their compounds?

- (a) Among the 3d-block elements, Sc has the highest density.  
 (b) The radii of atoms (Sc to Cu) increase from left to right.  
 (c)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$  is blue in colour whereas  $[\text{Zn}(\text{NH}_3)_4]^{2+}$  is colourless.  
 (d) The IUPAC name of  $\text{K}_2\text{NiCl}_4$  is dipotassium tetrachloronickelate(II).

32. Which statement/s is/are correct regarding the following molecule?




- (a) Atoms labelled P, Q, R and S lie on a straight line.  
 (b) Atoms labelled Q, R, S and T lie on a straight line.  
 (c) Atoms labelled R, S, T, U and V lie on the same plane.  
 (d) Atoms labelled R, S, T and U lie on a straight line.

33. Consider the equilibrium  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$  at 500 K.

Which of the following statement/s is/are correct for the above equilibrium?

- (a) Expression for the equilibrium constant for the equilibrium is,  $K_c = \frac{[\text{NH}_3(\text{g})]^2}{[\text{N}_2(\text{g})] \frac{1}{2} [\text{H}_2(\text{g})]^{\frac{3}{2}}}$
- (b) Expression for the equilibrium constant for the equilibrium is,  $K_c = \frac{[\text{N}_2(\text{g})]^{\frac{1}{2}} [\text{H}_2(\text{g})]^{\frac{3}{2}}}{[\text{NH}_3(\text{g})]^2}$
- (c) Expression for the equilibrium constant for the backward reaction is,  $K'_c = \frac{[\text{NH}_3(\text{g})]^2}{[\text{N}_2(\text{g})][\text{H}_2(\text{g})]^3}$
- (d) Expression for the equilibrium constant for the backward reaction is,  $K'_c = \frac{[\text{N}_2(\text{g})][\text{H}_2(\text{g})]^3}{[\text{NH}_3(\text{g})]^2}$

34. Which of the following statement/s regarding the reaction between compound **P** and HCl to form an alkyl halide is/are correct?



- (a) The major product is 2-chloro-2-methylbutane.  
 (b) A secondary carbocation is formed as an intermediate in this reaction.  
 (c) In one of the steps of the reaction, the HCl bond is cleaved to give a chlorine radical (Cl<sup>•</sup>).  
 (d) In one of the steps of the reaction, a nucleophile reacts with a carbocation.
35. A binary liquid mixture prepared by mixing two liquids in a closed evacuated container at a given temperature shows a negative deviation from Raoult's Law. Which of the following statement/s is/are correct for this system?
- (a) Total vapour pressure of the mixture is less than the expected total vapour pressure should it behave as an ideal mixture.  
 (b) Heat is released when the mixture is formed.  
 (c) Number of molecules in the vapour phase of the mixture is greater than the expected number of molecules should it behave as an ideal mixture.  
 (d) Heat is absorbed when the mixture is formed.
36. Which of the following statement/s is/are correct with regard to effects on natural cycles (e.g. N<sub>2</sub> cycle) present in our environment due to human activities?
- (a) Fossil fuel burning contributes to elevated levels of CO<sub>2</sub> in the atmosphere.  
 (b) CO<sub>2</sub> emission due to respiration of organisms causes elevated levels of CO<sub>2</sub> in the atmosphere.  
 (c) Industrial fixation of nitrogen causes increased amounts of nitrogen containing compounds in water and soil.  
 (d) Deforestation does not contribute to elevated levels of CO<sub>2</sub> in the atmosphere.
37. Which of the following statement/s is/are correct with regard to halogens, noble gases and their compounds?
- (a) Hypochlorous ion disproportionates rapidly in acidic solutions.  
 (b) Xe forms a series of compounds with F<sub>2</sub> gas, among which XeF<sub>4</sub> has a square planar geometry.  
 (c) Among the hydrogen halides, HF has the highest bond dissociation energy per mole.  
 (d) Boiling points of halogens increase down the group as a result of increasing strength of London forces.
38. Which of the following statement/s is/are correct regarding the Daniell cell when it operates at room temperature? ( $E_{cell}^{\circ} = +1.10 \text{ V}$ )
- (a) Net electron flow occurs from Zn to Cu.  
 (b) The equilibrium  $\text{Zn}^{2+}(\text{aq}) + 2\text{e} \rightleftharpoons \text{Zn}(\text{s})$  shifts to the right.  
 (c) A liquid-junction potential is created due to the presence of a salt-bridge.  
 (d) The equilibrium  $\text{Cu}^{2+}(\text{aq}) + 2\text{e} \rightleftharpoons \text{Cu}(\text{s})$  shifts to the right.
39. Which of the following statement/s is/are correct for ideal gases and real gases at constant temperature?
- (a) At very high pressures, the volume of a real gas is higher than that of an ideal gas.  
 (b) At high pressures, real gases tend to behave as ideal gases.  
 (c) At very high pressures, the volume of a real gas is lower than that of an ideal gas.  
 (d) At low pressures, real gases tend to behave as ideal gases.
40. Which of the following statement/s is/are correct regarding some industrial processes?
- (a) The first two steps involved in the manufacture of Na<sub>2</sub>CO<sub>3</sub> by Solvay Process are endothermic.  
 (b) Production of urea is carried out by a two step process using ammonia and carbon dioxide as the only raw materials.  
 (c) The first step involved in the manufacture of nitric acid by Ostwald method is the oxidation of NH<sub>3</sub> gas using O<sub>2</sub> in air in the presence of a catalyst to give NO<sub>2</sub> gas.  
 (d) High temperature and low pressure conditions are employed in the manufacture of NH<sub>3</sub> gas using Haber-Bosh process.

- In question Nos. 41 to 50, two statements are given in respect of each question. From the Table given below, select the response, out of the responses (1), (2), (3), (4) and (5), that best fits the two statements and mark appropriately on your answer sheet.

Response	First Statement	Second Statement
(1)	True	True, and correctly explains the first statement
(2)	True	True, but does <b>not</b> explain the first statement correctly
(3)	True	False
(4)	False	True
(5)	False	False

	First Statement	Second statement
41.	Among the oxides of Cr and Mn, CrO and MnO are acidic, while CrO <sub>3</sub> and Mn <sub>2</sub> O <sub>7</sub> are basic.	The acidic/basic nature of the oxides of Cr and Mn is dependant on the oxidation number of the metal.
42.	An acidic buffer solution can be prepared by mixing a weak acid HA(aq) with its sodium salt NaA(aq).	When OH <sup>-</sup> (aq) or H <sup>+</sup> (aq) ions are added to a buffer solution, the added amounts of OH <sup>-</sup> (aq) or H <sup>+</sup> (aq) ions are removed through the reactions; OH <sup>-</sup> (aq) + HA(aq) → A <sup>-</sup> (aq) + H <sub>2</sub> O(l) and H <sup>+</sup> (aq) + A <sup>-</sup> (aq) → HA(aq) respectively.
43.	Polyvinyl chloride (PVC) shows some resistance to catching fire.	Polyvinyl chloride (PVC) is a thermosetting polymer.
44.	At a given temperature and pressure the molar volumes of two different ideal gases are different from each other.	At 0 °C temperature and 1 atm pressure, the molar volume of an ideal gas is 22.4 dm <sup>3</sup> mol <sup>-1</sup> .
45.	All compounds having a C=C bond show diastereoisomerism.	Any two isomers which are not mirror images of each other are diastereoisomers.
46.	Hydrogenation of benzene is more difficult than hydrogenation of alkenes.	Addition of hydrogen to benzene results in the loss of aromatic stabilization.
47.	The reaction that takes place between SO <sub>3</sub> gas and water in the production of sulphuric acid is endothermic.	SO <sub>3</sub> gas reacts with concentrated H <sub>2</sub> SO <sub>4</sub> to give oleum.
48.	Reaction between ammonia and an alkylhalide gives a mixture of primary, secondary and tertiary amines and a quaternary ammonium salt.	Primary, secondary and tertiary amines can react as nucleophiles.
49.	If P + Q → R is a first order reaction with respect to the reactant P, the graph of rate against concentration of P gives a straight line passing through the origin.	Initial rate of a first order reaction is independent of the concentration of reactant(s).
50.	On a sunny day, strong photochemical smog can be seen in a city with heavy traffic congestion.	Photochemical smog is caused entirely by scattering of solar radiation by small particles and water droplets that are emitted by vehicle exhaust systems.

\*\*\*





සියලු ම හිමිකම් ඇවිරිණි / முழுப் பதிப்புரிமையுடையது / All Rights Reserved

පැරණි නිර්දේශය/பழைய பாடத்திட்டம் / Old Syllabus

ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව  
 திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம்  
 Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka  
 இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம் இலங்கைப் பரීட்சைத் திணைக்களம்

**OLD**

අධ්‍යයන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 2020

கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2020

General Certificate of Education (Adv. Level) Examination, 2020

රසායන විද්‍යාව II  
 இரசாயனவியல் II  
**Chemistry II**

**02 E II**


පැය තුනයි  
 மூன்று மணித்தியாலம்  
**Three hours**

අමතර කියවීමේ කාලය - මිනිත්තු 10 යි  
 மேலதிக வாசிப்பு நேரம் - 10 நிமிடங்கள்  
**Additional Reading Time - 10 minutes**

Use additional reading time to go through the question paper, select the questions and decide on the questions that you give priority in answering.

Index No. : .....

- \* A Periodic Table is provided on page 15.
- \* Use of calculators is not allowed.
- \* Universal gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
- \* Avogadro constant,  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$
- \* In answering this paper, you may represent alkyl groups in a condensed manner.



PART A – Structured Essay (pages 02 - 08)

- \* Answer all the questions on the question paper itself.
- \* Write your answer in the space provided for each question. Please note that the space provided is sufficient for the answer and that extensive answers are not expected.

PART B and PART C – Essay (pages 09 - 14)

- \* Answer four questions selecting two questions from each part. Use the papers supplied for this purpose.
- \* At the end of the time allotted for this paper, tie the answers to the three Parts A, B and C together so that Part A is on top and hand them over to the Supervisor.
- \* You are permitted to remove only Parts B and C of the question paper from the Examination Hall.

For Examiner's Use Only

Part	Question No.	Marks
A	1	
	2	
	3	
	4	
B	5	
	6	
	7	
C	8	
	9	
	10	
Total		

Total

In Numbers	
In Letters	

Code Numbers

Marking Examiner 1	
Marking Examiner 2	
Checked by :	
Supervised by :	

## PART A – STRUCTURED ESSAY

Answer all four questions on this paper itself. (Each question carries 10 marks.)

Do not write in this column.

1. (a) Write the answers to the questions given below on the dotted lines.

(i) Of the three ions  $\text{Na}^+$ ,  $\text{Mg}^{2+}$  and  $\text{F}^-$ , which one has the **smallest** ionic radius? .....

(ii) Of the three elements C, N and O, which one has the **highest** second ionization energy? .....

(iii) Of the three compounds  $\text{H}_2\text{O}$ ,  $\text{HOCl}$  and  $\text{OF}_2$ , which one has the **most** electronegative oxygen atom? .....


(iv) Of the three elements Be, C and N, which one will liberate energy when an electron is added to its atom [ $\text{Y}(\text{g}) + \text{e} \rightarrow \text{Y}^-(\text{g})$ ;  $\text{Y} = \text{Be}, \text{C}, \text{N}$ ] in the gaseous state? .....

(v) Of the three ionic compounds  $\text{NaF}$ ,  $\text{KF}$  and  $\text{KBr}$ , which one has the **highest** solubility in water? .....

(vi) Of the three compounds  $\text{HCHO}$ ,  $\text{CH}_3\text{F}$  and  $\text{H}_2\text{O}_2$ , which one has the **strongest** intermolecular forces? .....



(2.4 marks)

(b) (i) Draw the **most** acceptable Lewis structure for the ion,  $\text{N}_2\text{O}_3^{2-}$ . Its skeleton is given below.



(ii) Draw **three** more Lewis structures (resonance structures) for this ion. Indicate the relative stabilities of the structures drawn by you, when compared with the most acceptable structure drawn in (i) above, by writing 'less stable' or 'unstable' under these structures.

(iii) Complete the given table based on the Lewis structure and its labelled skeleton given below.



	N <sup>1</sup>	N <sup>2</sup>	O <sup>3</sup>	C <sup>4</sup>
VSEPR pairs around the atom				
electron pair geometry around the atom				
shape around the atom				
hybridization of the atom				

[see page three

Do not  
write  
in this  
column.

- Parts (iv) to (vii) are based on the Lewis structure given in part (iii) above. Labelling of atoms is as in part (iii).

(iv) Identify the atomic/hybrid orbitals involved in the formation of  $\sigma$  bonds between the two atoms given below.

- |                                     |                      |                      |
|-------------------------------------|----------------------|----------------------|
| I. Cl—N <sup>1</sup>                | Cl .....             | N <sup>1</sup> ..... |
| II. N <sup>1</sup> —O               | N <sup>1</sup> ..... | O .....              |
| III. N <sup>1</sup> —N <sup>2</sup> | N <sup>1</sup> ..... | N <sup>2</sup> ..... |
| IV. N <sup>2</sup> —O <sup>3</sup>  | N <sup>2</sup> ..... | O <sup>3</sup> ..... |
| V. O <sup>3</sup> —C <sup>4</sup>   | O <sup>3</sup> ..... | C <sup>4</sup> ..... |
| VI. C <sup>4</sup> —N               | C <sup>4</sup> ..... | N .....              |

(v) Identify the atomic orbitals involved in the formation of  $\pi$  bonds between the two atoms given below.

- |                                   |                      |                      |
|-----------------------------------|----------------------|----------------------|
| I. N <sup>1</sup> —N <sup>2</sup> | N <sup>1</sup> ..... | N <sup>2</sup> ..... |
| II. C <sup>4</sup> —N             | C <sup>4</sup> ..... | N .....              |
|                                   | C <sup>4</sup> ..... | N .....              |

(vi) State the approximate bond angles around N<sup>1</sup>, N<sup>2</sup>, O<sup>3</sup> and C<sup>4</sup> atoms.

N<sup>1</sup> ....., N<sup>2</sup> ....., O<sup>3</sup> ....., C<sup>4</sup> .....

(vii) Arrange the atoms N<sup>1</sup>, N<sup>2</sup>, O<sup>3</sup> and C<sup>4</sup> in the **increasing** order of electronegativity.

..... < ..... < ..... < ..... (5.6 marks)

(c) Consider the following information.

I. The atoms **A** and **B** combine to form a heterodiatomic molecule **AB** that has a  $\sigma$  bond. This is represented as **A—B**.

II. The electronegativity of **A** is less than that of **B** ( $X_A < X_B$ ).  
X = electronegativity of the atom

III. The inter-nuclear distance between **A** and **B** atoms ( $d_{A-B}$ ) of the **AB** molecule is given by the following equation.

$$d_{A-B} = r_A + r_B - c(X_B - X_A)$$

r = atomic radius, c = 9 pm

**Note:** d and r are measured in picometres (pm). (1 pm = 10<sup>-12</sup> m)

Based on the above information, answer the following questions.

(i) What is the name used to identify the type of  $\sigma$  bond between **A** and **B**?

.....

(ii) Show how fractional charges ( $\delta+$  and  $\delta-$ ) are located in the molecule **AB**.

.....

(iii) Write the equation to calculate the dipole moment ( $\mu$ ) of molecule **AB** and show its direction.

- (iv) Calculate the percentage of ionic character of the H-F bond in the HF molecule using the data given below.

Inter-nuclear distance of  $H_2$  ( $d_{H-H}$ ) = 74 pm

Electronegativity of F = 4.0

Inter-nuclear distance of  $F_2$  ( $d_{F-F}$ ) = 144 pm

Dipole moment of HF =  $6.0 \times 10^{-30}$  C m

Electronegativity of H = 2.1

Charge of an electron =  $1.6 \times 10^{-19}$  C

Do not write in this column.

(2.0 marks)

100

2. (a) A, B, C and D are chlorides of *p*-block elements. The elements A, B and C have atomic numbers less than 20 whereas in D it is greater than 20 ( $20 < Z_D < 55$ ). A description of the products ( $P_1$ – $P_8$ ) formed when A is reacted with a limited amount of water and B, C and D are reacted with excess water are given below.

Compound	Description of products
A	$P_1$ a very weakly acidic solid
	$P_2$ a strong monobasic acid
B	$P_3$ a gas that turns red litmus blue
	$P_4$ a compound with bleaching properties
C	$P_5$ a tribasic acid
	$P_6$ a strong monobasic acid
D	$P_7$ a white precipitate
	$P_8$ a strong monobasic acid

- (i) Identify A, B, C and D (give the chemical formulae).

A: ..... B: ..... C: ..... D: .....

- (ii) Give balanced chemical equations for the reactions of A, B, C and D with water to give products  $P_1$  to  $P_8$ .

.....  
 .....  
 .....  
 .....

Do not  
write  
in this  
column.

(iii) Write balanced chemical equations for the following reactions.

I.  $P_1$  with NaOH(aq)

.....

II.  $P_3$  with Mg

.....

III.  $P_8$  with Al

.....

(5.0 marks)

(b) A student is provided with bottles labelled **P, Q, R, S, T** and **U** containing aqueous solutions of  $Al_2(SO_4)_3$ ,  $H_2SO_4$ ,  $Na_2S_2O_3$ ,  $BaCl_2$ ,  $Pb(Ac)_2$  and  $KOH$  (**not in order**). Some useful observations for their identification on mixing two solutions at a time are given below.

(Ac - Acetate ion)

	Solutions mixed	Observations
I	T + R	a clear colourless solution
II	P + R	a white precipitate
III	T + S	a gelatinous white precipitate
IV	U + R	a white precipitate
V	P + Q	a white precipitate, turns black on heating
VI	P + U	a white precipitate, dissolves on heating

(i) Identify **P** to **U**.**P:** .....**Q:** .....**R:** .....**S:** .....**T:** .....**U:** .....(ii) Give balanced chemical equations for each of the reactions **I** to **VI**.**I:** .....**II:** .....**III:** .....**IV:** .....**V:** formation of white precipitate: .....

turning black on heating: .....

**VI:** .....

(Note: indicate precipitates as ↓)

(5.0 marks)

3. (a) A saturated aqueous solution of a sparingly soluble salt  $AB_2(s)$  was prepared by stirring an excess amount of  $AB_2(s)$  in  $1.0 \text{ dm}^3$  of distilled water at  $25^\circ\text{C}$ . The amount of  $A^{2+}(aq)$  ions present in this saturated aqueous solution was found to be  $2.0 \times 10^{-3} \text{ mol}$ .

(i) Write the equilibrium related to the dissolution of  $AB_2(s)$  in the above system at  $25^\circ\text{C}$ .

.....

(ii) Write the expression for the equilibrium constant for the equilibrium written in (i) above at  $25^\circ\text{C}$ .

.....

.....

100

- (iv) Calculate the percentage of ionic character of the H-F bond in the HF molecule using the data given below.

Inter-nuclear distance of $H_2$ ( $d_{H-H}$ ) = 74 pm	Electronegativity of F = 4.0
Inter-nuclear distance of $F_2$ ( $d_{F-F}$ ) = 144 pm	Dipole moment of HF = $6.0 \times 10^{-30}$ C m
Electronegativity of H = 2.1	Charge of an electron = $1.6 \times 10^{-19}$ C

Do not write in this column.

(2.0 marks)

2. (a) A, B, C and D are chlorides of *p*-block elements. The elements A, B and C have atomic numbers less than 20 whereas in D it is greater than 20 ( $20 < Z_D < 55$ ). A description of the products ( $P_1 - P_8$ ) formed when A is reacted with a limited amount of water and B, C and D are reacted with excess water are given below.

Compound	Description of products	
A	$P_1$	a very weakly acidic solid
	$P_2$	a strong monobasic acid
B	$P_3$	a gas that turns red litmus blue
	$P_4$	a compound with bleaching properties
C	$P_5$	a tribasic acid
	$P_6$	a strong monobasic acid
D	$P_7$	a white precipitate
	$P_8$	a strong monobasic acid

- (i) Identify A, B, C and D (give the chemical formulae).

A: ..... B: ..... C: ..... D: .....

- (ii) Give balanced chemical equations for the reactions of A, B, C and D with water to give products  $P_1$  to  $P_8$ .

.....

.....

.....

.....

PAPERMASTER.LK

100

Do not  
write  
in this  
column.




4. (a) **A**, **B**, **C** and **D** are structural isomers having the molecular formula  $C_6H_{10}$ . None of them show optical isomerism. All four isomers, **A**, **B**, **C** and **D** when treated with  $HgSO_4/dil. H_2SO_4$  give products which react with 2,4-dinitrophenylhydrazine (2,4-DNP) to give coloured precipitates.

Only **A** gives a precipitate with ammonical  $AgNO_3$ . **A** has only one position isomer, which is **B**.

**B** is a chain isomer of **C**. **C** reacts with  $HgSO_4/dil. H_2SO_4$  to give two products **E** and **F**.


**D** reacts with  $HgSO_4/dil. H_2SO_4$  to give only one product, which is **E**.

(i) Draw the structures of **A**, **B**, **C**, **D**, **E** and **F** in the boxes given below.




**A****B****C****D****E****F**

(ii) Which of the compounds **A**, **B**, **C** and **D** gives a product that does not show diastereoisomerism when reacted separately with  $H_2 / Pd-BaSO_4 /$  quinoline?

(iii) Draw, in the box given below, the structure of the product **G** obtained when **A** is reacted with excess  $HBr$ .

**G**

(iv) Draw the structures of products **X** and **Y** obtained in the following reactions of **E**, in the appropriate boxes.

**X****Y**

Name a test to distinguish between **X** and **Y**.

.....



(6.0 marks)




- (b) (i) Complete the following three reaction sequences by drawing structures of compounds K, L and M and giving the reagents/catalysts P, Q and R in the boxes given below.

Do not write in this column.

Sequence 1:





P




K


Sequence 2:



L





Q



M

Sequence 3:



R

(3.0 marks)

- (ii) Selecting from the reactions I–VI, give one (01) example for each of the following types of reactions.

Nucleophilic addition .....

Nucleophilic substitution .....

(1.0 marks)

\*\*

100

## පැරණි නිර්දේශය/பழைய பாடத்திட்டம்/ Old Syllabus

ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව  
 இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம்  
 Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka Department of Examinations, Sri Lanka  
 இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம் இலங்கைப் பரட்சைத் திணைக்களம்  
 Department of Examinations, Sri Lanka

**OLD**

අධ්‍යයන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 2020  
 கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர) பரீட்சை, 2020  
 General Certificate of Education (Adv. Level) Examination, 2020

රසායන විද්‍යාව II  
 இரசாயனவியல் II  
**Chemistry II**

**02 E II**


\* Universal gas constant  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$

\* Avogadro constant  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

## PART B – ESSAY

Answer two questions only. (Each question carries 15 marks.)

5. (a) A compound  $\text{XY}_2\text{Z}_2(\text{g})$  undergoes dissociation when heated to temperatures above 300 K as given below.




A sample of 7.5 g of  $\text{XY}_2\text{Z}_2(\text{g})$  was placed in an evacuated 1.00 dm<sup>3</sup> rigid-closed container and the temperature was raised to 480 K.

Molar mass of  $\text{XY}_2\text{Z}_2(\text{g})$  is 150 g mol<sup>-1</sup>. Use the approximate value of 4000 J mol<sup>-1</sup> for  $RT$  at 480 K. Assume ideal gas behaviour for all gases.

- Calculate the number of moles of  $\text{XY}_2\text{Z}_2(\text{g})$  in the container before dissociation.
  - When the above system reaches equilibrium at 480 K, the total number of moles in the container was found to be  $7.5 \times 10^{-2}$  mol. Calculate the number of moles of  $\text{XY}_2\text{Z}_2(\text{g})$ ,  $\text{XY}_2(\text{g})$  and  $\text{Z}_2(\text{g})$  in the equilibrium mixture at 480 K.
  - Calculate the equilibrium constant  $K_c$  for the above reaction at 480 K.
  - Calculate  $K_p$  for the equilibrium at 480 K. (7.5 marks)
- (b) For the reaction  $\text{XY}_2\text{Z}_2(\text{g}) \rightarrow \text{XY}_2(\text{g}) + \text{Z}_2(\text{g})$  described in (a), Gibbs free energies ( $G$ ) at 480 K for  $\text{XY}_2\text{Z}_2(\text{g})$ ,  $\text{XY}_2(\text{g})$  and  $\text{Z}_2(\text{g})$  are  $-60 \text{ kJ mol}^{-1}$ ,  $-76 \text{ kJ mol}^{-1}$  and  $-30 \text{ kJ mol}^{-1}$ , respectively.
- Calculate  $\Delta G$  (in kJ mol<sup>-1</sup>) for the reaction at 480 K.
  - The magnitude of  $\Delta S$  of the above reaction is  $150 \text{ J K}^{-1} \text{ mol}^{-1}$  at 480 K. Calculate  $\Delta H$  for the reaction at 480 K by using the appropriate sign (- or +) of  $\Delta S$ .
  - By using the sign (- or +) of  $\Delta H$  obtained in (ii), explain whether this reaction is exothermic or endothermic.
  - Deduce the enthalpy difference for the formation of  $\text{XY}_2\text{Z}_2(\text{g})$  from  $\text{XY}_2(\text{g})$  and  $\text{Z}_2(\text{g})$  at 480 K.
  - If the bond enthalpy of the X-Z bond in  $\text{XY}_2\text{Z}_2(\text{g})$  is  $+250 \text{ kJ mol}^{-1}$ , calculate the bond enthalpy of the Z-Z bond.

(Assume that  $\text{XY}_2\text{Z}_2(\text{g})$  has the structure  $\text{Z}-\text{X}-\text{Z}$ )



- If liquid  $\text{XY}_2\text{Z}_2$  is used instead of gaseous  $\text{XY}_2\text{Z}_2$ , giving reasons, explain whether the value of  $\Delta H$  obtained for the reaction  $\text{XY}_2\text{Z}_2(\text{l}) \rightarrow \text{XY}_2(\text{g}) + \text{Z}_2(\text{g})$  is equal to, or higher or lower than  $\Delta H$  obtained in (ii). (7.5 marks)

6. (a) Consider the reaction given below occurring in a closed container at a given temperature  $T$ .



- (i) Write **three** expressions for the rate of reaction relevant to each of the compounds appearing in the reaction.
- (ii) This reaction was carried out at temperature  $T$  with an initial concentration of  $0.10 \text{ mol dm}^{-3}$  of  $\text{N}_2\text{O}_5(\text{g})$ . It was found that 40% of the initial amount was decomposed after a period of 400 s.
- Calculate the average rate of decomposition of  $\text{N}_2\text{O}_5(\text{g})$  in this time interval.
  - Calculate average rates of formation of  $\text{NO}_2(\text{g})$  and  $\text{O}_2(\text{g})$ .
- (iii) In another experiment, initial rates were measured for this reaction at 300 K and the results are given below.


$[\text{N}_2\text{O}_5(\text{g})] / \text{mol dm}^{-3}$	0.01	0.02	0.03
Initial rate / $\text{mol dm}^{-3} \text{ s}^{-1}$	$6.930 \times 10^{-5}$	$1.386 \times 10^{-4}$	$2.079 \times 10^{-4}$

Derive the rate law for the reaction at 300 K.

- (iv) Another experiment was carried out at 300 K with an initial concentration of  $0.64 \text{ mol dm}^{-3}$  of  $\text{N}_2\text{O}_5(\text{g})$ . It was found that the concentration of  $\text{N}_2\text{O}_5(\text{g})$  which remained after a period of 500 s was  $2.0 \times 10^{-2} \text{ mol dm}^{-3}$ .

- Calculate the half-life ( $t_{1/2}$ ) of the reaction at 300 K.
- Calculate the rate constant of the reaction at 300 K.

- (v) This reaction proceeds through a mechanism involving the following elementary steps.



Show that the above mechanism is consistent with the rate law of the reaction. **(8.0 marks)**

- (b) An ideal binary-liquid mixture was prepared by mixing two liquids of **A** and **B** in a closed evacuated container at temperature  $T$ . After establishing the equilibrium at temperature  $T$ , partial pressures of **A** and **B** in the vapour phase are  $P_A$  and  $P_B$ , respectively. At temperature  $T$ , the saturated vapour pressures of **A** and **B** are  $P_A^\circ$  and  $P_B^\circ$ , respectively. Mole fractions of **A** and **B** in solution are  $X_A$  and  $X_B$ , respectively.

- (i) Show that  $P_A = P_A^\circ X_A$

(Consider that the rates of vaporization and condensation are equal at equilibrium.)


- (ii) In the above system at 300 K, the total pressure was  $5.0 \times 10^4 \text{ Pa}$ . The saturated vapour pressures of pure **A** and **B** at 300 K, are  $7.0 \times 10^4 \text{ Pa}$  and  $3.0 \times 10^4 \text{ Pa}$ , respectively.

- Calculate the mole fraction of **A** in the liquid phase of the equilibrium mixture.
- Calculate the vapour pressure of **A** in the equilibrium mixture.

**(7.0 marks)**

7. (a) (i) Given below is a reaction scheme for the synthesis of compound G using  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$  as the only organic starting compound.

Complete the reaction scheme by drawing the structures of compounds A, B, C, D, E and F and writing the appropriate reagents for steps 1 – 7, selected only from those given in the list.




**List of Reagents**

HBr,  $\text{PBr}_3$ , pyridiniumchlorochromate (PCC),  
Mg / dry ether, KCN, conc.  $\text{H}_2\text{SO}_4$ , dil.  $\text{H}_2\text{SO}_4$

(5.2 marks)

- (ii) Consider the following series of reactions.


Draw the structures of compounds G, H and K. Give the reagents X, Y and Z.



Note that K gives benzyl alcohol ( $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$ ) when reacted with  $\text{NaNO}_2 / \text{dil. HCl}$ .


(2.4 marks)

- (b) (i) Show how the following conversion could be carried out in **not more than three steps**.



(2.0 marks)

- (ii) Consider the following reaction.



Identify the chemical substances P and Q necessary to carry out this reaction.

Write the mechanism of this reaction.

(2.0 marks)


- (c) (i) Explain why phenol is more reactive in electrophilic substitution reactions than benzene, by considering their resonance hybrids.
- (ii) Illustrate the difference in reactivity between phenol and benzene as given in (i) above by means of a suitable reaction.
- (iii) Draw the structure(s) of product(s) you described in the reaction in (ii) above.

(3.4 marks)

## PART C – ESSAY

Answer two questions only. (Each question carries 15 marks.)

8. (a) (i) Write the chemical formulae of the substances A – Q given in the flow chart below. (Note: Chemical equations and reasons are not expected for the identification of substances A – Q.) The symbols given in the box (dash lines) are used to represent solids, precipitates, solutions and gases.



- (ii) Write the complete electronic configuration of A.
- (iii) State the function of E in the conversion of D to F. Give the relevant balanced chemical equations for the stated function. (7.5 marks)
- (b) The solid X contains only  $\text{Cu}_2\text{S}$  and  $\text{CuS}$ . The following procedure was used to determine the percentage of  $\text{Cu}_2\text{S}$  in X.

**Procedure**

A 1.00 g portion of solid X was treated with  $100.00 \text{ cm}^3$  of  $0.16 \text{ mol dm}^{-3} \text{ KMnO}_4$  in dilute  $\text{H}_2\text{SO}_4$  medium. This reaction gave  $\text{Mn}^{2+}$ ,  $\text{Cu}^{2+}$  and  $\text{SO}_4^{2-}$  as products. Thereafter, the excess  $\text{KMnO}_4$  in this solution was titrated with  $0.15 \text{ mol dm}^{-3} \text{ Fe}^{2+}$  solution. The volume required for the titration was  $35.00 \text{ cm}^3$ .

- (i) Write the balanced ionic equations for the reactions taking place in the above procedure.
- (ii) Based on the answers to (i) above, determine the molar ratio between,
- $\text{Cu}_2\text{S}$  and  $\text{KMnO}_4$
  - $\text{CuS}$  and  $\text{KMnO}_4$
  - $\text{Fe}^{2+}$  and  $\text{KMnO}_4$
- (iii) Calculate the percentage by weight of  $\text{Cu}_2\text{S}$  in X. (Cu = 63.5, S = 32) (7.5 marks)

[see page thirteen]

9. (a) The following questions are based on the production of NaOH using the membrane cell method.
- (i) State the raw material used.
  - (ii) Draw and label fully the membrane cell used in the manufacturing process.
  - (iii) Briefly explain the manufacturing process using balanced chemical equations where applicable.
  - (iv) Give **three** uses of NaOH.
  - (v) Give **two** uses of each by-product formed in the manufacturing process.


(7.5 marks)

- (b) Currently, global warming due to change in greenhouse effect is significantly greater than that before the industrial revolution.

- (i) Explain briefly what is meant by greenhouse effect.
- (ii) State **two main** natural gases that contribute to global warming.
- (iii) Briefly explain the direct relationship between human impact on carbon cycle and global warming.
- (iv) Explain briefly how microorganisms contribute to the release of the gases you stated in (ii).
- (v) In addition to the gases you stated in (ii), name a class of synthetic volatile compounds that directly contribute to global warming and draw the structure of a selected compound from this class.
- (vi) Name **five** effects caused by global warming on global water.
- (vii) The slow down of industrial activities due to the Covid-19 pandemic temporarily eased the global environmental issues in many countries. Justify this statement by using **three** main global environmental issues you have learnt.

(7.5 marks)

10. (a) A coloured complex ion **P** is formed when the salt  $M(NO_3)_n$  is dissolved in distilled water. **M** is a transition element belonging to the 3d block. **P** undergoes the following reactions.



- (i) Identify the metal **M**. Give the oxidation state of **M** in complex ion **P**.
  - (ii) Give the value of  $n$  in  $M(NO_3)_n$ .
  - (iii) Write the complete electronic configuration of **M** in complex ion **P**.
  - (iv) Write the chemical formulae of **P, Q, R, S, T** and **U**.
  - (v) Give the IUPAC names of **P, R** and **S**.
  - (vi) What is the colour of **R**?
  - (vii) What would you expect to observe when aqueous solutions of the following compounds are treated with **U**?
    - I.  $MnCl_2$
    - II.  $ZnCl_2$
  - (viii) When aq.  $NH_3$  is added to a solution of a salt of a transition metal, a yellow-brown complex ion **V** is formed.
    - I. Identify **V**.
    - II. Give the IUPAC name of **V**.
  - (ix) Briefly describe a method with the aid of balanced chemical equations for determining the concentration of  $M^{n+}$  present in an aqueous solution, using the following chemicals.  
 $KI, Na_2S_2O_3$  and starch. (7.5 marks)
- (b) (i) To compare the properties of Electrolytic and Galvanic cells, copy and complete the following table using the given terms.

Terms: anode, cathode, positive, negative, spontaneous, non-spontaneous.

	Electrolytic cell	Galvanic cell
A. Oxidation half-reaction takes place at		
B. Reduction half-reaction takes place at		
C. Sign of $E_{cell}^{\circ}$		
D. Electron flow	From ..... to .....	From ..... to .....
E. Spontaneity of the cell reaction		

- (ii) An electrochemical cell was constructed at 300 K by using a  $Zn(s)$  anode, an aqueous alkaline electrolyte and a porous Pt cathode which facilitates the collection of oxygen  $O_2(g)$  from air as shown below. As the cell operates  $ZnO(s)$  is produced.

You are given that

$$E_{ZnO(s)|Zn(s)|OH^-(aq)}^{\circ} = -1.31 \text{ V and } E_{O_2(g)|OH^-(aq)}^{\circ} = +0.34 \text{ V}$$

$$Zn = 65 \text{ g mol}^{-1}, O = 16 \text{ g mol}^{-1} \text{ and}$$

$$1 F = 96,500 \text{ C}$$

- I. Write the half-reactions occurring at anode and cathode.

- II. Write the overall cell reaction.

- III. Calculate the cell potential  $E_{cell}^{\circ}$  at 300 K.

- IV. State the direction of migration of  $OH^-(aq)$  ions between the electrodes.


- V. When the cell operates for a period of 800 s at 300 K, 2 mol of  $O_2(g)$  are consumed.

- A. Calculate the number of moles of electrons passing through the cell.

- B. Calculate the mass of  $ZnO(s)$  formed.

- C. Calculate the current passing through the cell.

(7.5 marks)



The Periodic Table

1	1 <b>H</b>																	2 <b>He</b>
2	3 <b>Li</b>	4 <b>Be</b>											5 <b>B</b>	6 <b>C</b>	7 <b>N</b>	8 <b>O</b>	9 <b>F</b>	10 <b>Ne</b>
3	11 <b>Na</b>	12 <b>Mg</b>											13 <b>Al</b>	14 <b>Si</b>	15 <b>P</b>	16 <b>S</b>	17 <b>Cl</b>	18 <b>Ar</b>
4	19 <b>K</b>	20 <b>Ca</b>	21 <b>Sc</b>	22 <b>Ti</b>	23 <b>V</b>	24 <b>Cr</b>	25 <b>Mn</b>	26 <b>Fe</b>	27 <b>Co</b>	28 <b>Ni</b>	29 <b>Cu</b>	30 <b>Zn</b>	31 <b>Ga</b>	32 <b>Ge</b>	33 <b>As</b>	34 <b>Se</b>	35 <b>Br</b>	36 <b>Kr</b>
5	37 <b>Rb</b>	38 <b>Sr</b>	39 <b>Y</b>	40 <b>Zr</b>	41 <b>Nb</b>	42 <b>Mo</b>	43 <b>Tc</b>	44 <b>Ru</b>	45 <b>Rh</b>	46 <b>Pd</b>	47 <b>Ag</b>	48 <b>Cd</b>	49 <b>In</b>	50 <b>Sn</b>	51 <b>Sb</b>	52 <b>Te</b>	53 <b>I</b>	54 <b>Xe</b>
6	55 <b>Cs</b>	56 <b>Ba</b>	57 <b>La</b>	72 <b>Hf</b>	73 <b>Ta</b>	74 <b>W</b>	75 <b>Re</b>	76 <b>Os</b>	77 <b>Ir</b>	78 <b>Pt</b>	79 <b>Au</b>	80 <b>Hg</b>	81 <b>Tl</b>	82 <b>Pb</b>	83 <b>Bi</b>	84 <b>Po</b>	85 <b>At</b>	86 <b>Rn</b>
7	87 <b>Fr</b>	88 <b>Ra</b>	89 <b>Ac</b>	104 <b>Rf</b>	105 <b>Db</b>	106 <b>Sg</b>	107 <b>Bh</b>	108 <b>Hs</b>	109 <b>Mt</b>	110 <b>Ds</b>	111 <b>Rg</b>	112 <b>Cn</b>	113 <b>Nh</b>	114 <b>Fl</b>	115 <b>Mc</b>	116 <b>Lv</b>	117 <b>Ts</b>	118 <b>Og</b>

57 <b>La</b>	58 <b>Ce</b>	59 <b>Pr</b>	60 <b>Nd</b>	61 <b>Pm</b>	62 <b>Sm</b>	63 <b>Eu</b>	64 <b>Gd</b>	65 <b>Tb</b>	66 <b>Dy</b>	67 <b>Ho</b>	68 <b>Er</b>	69 <b>Tm</b>	70 <b>Yb</b>	71 <b>Lu</b>
89 <b>Ac</b>	90 <b>Th</b>	91 <b>Pa</b>	92 <b>U</b>	93 <b>Np</b>	94 <b>Pu</b>	95 <b>Am</b>	96 <b>Cm</b>	97 <b>Bk</b>	98 <b>Cf</b>	99 <b>Es</b>	100 <b>Fm</b>	101 <b>Md</b>	102 <b>No</b>	103 <b>Lr</b>